

STRONG-COUPLING EXPANSION FOR THE HUBBARD MODEL IN ARBITRARY DIMENSION USING SLAVE BOSONS

P.J.H. Denteneer*

Instituut–Lorentz, University of Leiden,
P. O. Box 9506, 2300 RA Leiden, The Netherlands

Abstract

A strong-coupling expansion for the antiferromagnetic phase of the Hubbard model is derived in the framework of the slave-boson mean-field approximation. The expansion can be obtained in terms of moments of the density of states of freely hopping electrons on a lattice, which in turn are obtained for hypercubic lattices in arbitrary dimension. The expansion is given for the case of half-filling and for the energy up to fifth order in the ratio of hopping integral t over on-site interaction U , but can straightforwardly be generalized to the non-half-filled case and be extended to higher orders in t/U . For the energy the expansion is found to have an accuracy of better than 1% for $U/t \geq 8$. A comparison is given with an earlier perturbation expansion based on the Linear Spin Wave approximation and with a similar expansion based on the Hartree-Fock approximation. The case of an infinite number of spatial dimensions is discussed.

PACS: 75.10.Lp, 02.30.Mv, 71.27.+a

Short title: Strong-coupling expansion for the Hubbard model

INLO-PUB 23/95; January 1996; to be published in Physical Review B.

*e-mail address: pjhdent@rulkol.leidenuniv.nl

1 Introduction

If the physics of strongly correlated fermions is to be described by the Hubbard model [1], electronic structure calculations for the cuprates indicate that the relevant parameter regime is where the on-site interaction U is comparable to or larger than the bandwidth W of freely hopping electrons [2]. A natural strategy to try to describe this regime is to approach it from the two limiting cases of weak-coupling ($U \ll W$) and strong-coupling ($U \gg W$). The simplest mean-field approximation for the Hubbard model, the Hartree-Fock approximation (HFA), is at first sight a weak-coupling approximation although, with some care, also a reasonable description of the strong-coupling regime can be given [3]–[6]. Strong-coupling approaches have mainly been devoted to the one-dimensional case (see e.g. [7]) and often take the limit $U \rightarrow \infty$ [8]. A somewhat more sophisticated mean-field approximation, namely that based on the slave-boson formulation due to Kotliar and Ruckenstein [9], is in principle not restricted to weak or strong coupling. In fact, it was shown that this so-called slave-boson mean-field (SBMF) approximation is equivalent to the Gutzwiller approximation to the Gutzwiller wave function [10], the latter of which improves upon the HFA especially in the *intermediate*-coupling regime. This SBFM approximation then allows to approach the interesting intermediate-coupling regime from the strong-coupling side. Such an approximate strong-coupling approach can be particularly helpful since more rigorous Quantum Monte Carlo calculations become increasingly cumbersome for stronger coupling [11].

In a previous work [12], we compared the HFA and SBFM approximation for various simple magnetic phases, as well as computed the effective hopping and spin stiffness for the *two-dimensional* (one-band) Hubbard model. In this paper, I derive a large- U expansion within the SBFM approximation for the Hubbard model in *arbitrary* dimension. Here, I restrict myself to the antiferromagnetic phase at half-filling, but the extension to doping the antiferromagnet with electrons or holes is straightforward (though tedious).

In Section 2, the slave-boson mean-field approximation is briefly introduced and the resulting set of integral equations for the antiferromagnetic phase is given. A large- U expansion is derived in Section 3 in terms of moments of the density of states for freely hopping electrons on a lattice, and compared to earlier numerical and analytical work in the literature. In Section 4, the expansion is made explicit for hypercubic lattices of arbitrary dimension. Finally, I discuss the convergence of the series and possible extensions. The Appendix contains useful formulae for the moments of the above mentioned density of states.

2 Slave-Boson Mean-Field Approximation

The Hubbard Hamiltonian is [1]:

$$\mathcal{H}_h = -t \sum_{j,\delta,\sigma} c_{j+\delta,\sigma}^\dagger c_{j\sigma} + U \sum_j n_{j\uparrow} n_{j\downarrow}, \quad (1)$$

where $c_{j\sigma}$ is the annihilation operator for an electron at site j with spin σ . Neighboring sites of site j are denoted by $j + \delta$. t is the hopping integral, U the on-site

interaction between electrons of opposite spin, and $n_{j\sigma} = c_{j\sigma}^\dagger c_{j\sigma}$ is the occupation number operator. The slave-boson approach of Kotliar and Ruckenstein consists of introducing bosons for each of the (four) possible electron occupancies of a site. The electron creation and annihilation operators are then modified such that always the (one) boson corresponding to the electron occupancy is present at each site. In a functional-integral description of the new Hamiltonian the slave-boson mean-field (SBMF) approximation is when the Bose fields are independent of (imaginary-) time. Different phases can be considered by assuming different forms of position dependence of the Bose fields. For more details see Refs.[9, 12]. The SBFM approximation is an improvement over the Hartree-Fock approximation since it takes some local correlations into account; in particular, the density of doubly occupied sites is an independent parameter to be optimized.

The set of equations describing the antiferromagnetic (AF) phase within SBFM is given by [12]:

$$m_s = 2\bar{\lambda} \int_{\bar{\mu}}^{\infty} d\varepsilon \frac{\mathcal{N}(\varepsilon)}{(\varepsilon^2 + \bar{\lambda}^2)^{1/2}} \quad (2)$$

$$\bar{\lambda} = \frac{2}{q_s} [q_s]_{m_s} \int_{\bar{\mu}}^{\infty} d\varepsilon \frac{\mathcal{N}(\varepsilon) \varepsilon^2}{(\varepsilon^2 + \bar{\lambda}^2)^{1/2}} \quad (3)$$

$$U = \frac{[q_s]_d}{d} \int_{\bar{\mu}}^{\infty} d\varepsilon \frac{\mathcal{N}(\varepsilon) \varepsilon^2}{(\varepsilon^2 + \bar{\lambda}^2)^{1/2}} \quad (4)$$

$$n = 2 \int_{\bar{\mu}}^{\infty} d\varepsilon \mathcal{N}(\varepsilon) , \quad (5)$$

where $\bar{\lambda}$ is an internal (“renormalized gap”) parameter; $\bar{\mu}$ an effective chemical potential; and the band renormalization q_s is a function of density n , sublattice magnetization m_s , and d (d^2 is the density of doubly occupied sites), and given by:

$$q_s(n, m_s, d) = z(n, m_s, d) z(n, -m_s, d) , \quad (6)$$

where

$$z(n, m_s, d) = \frac{\sqrt{(1 - n + d^2)(n + m_s - 2d^2)} + d\sqrt{n - m_s - 2d^2}}{\sqrt{(n + m_s) \left(1 - \frac{n + m_s}{2}\right)}} . \quad (7)$$

$[q_s]_\alpha$ is the derivative of q_s with respect to α (explicit expressions can be found in Ref.[12]) and $\mathcal{N}(\varepsilon)$ is the density of states (DOS) of freely hopping electrons. The dimension-dependence resides solely in $\mathcal{N}(\varepsilon)$ and since this DOS applies to a non-interacting system, I will be able to derive results for arbitrary dimension in Section 4 (see also the appendix). Note that the above set of equations reduces to the familiar antiferromagnetic or spin-density wave (SDW) solution in the HFA if $q_s = 1$ [3]. In that case, one has one self-consistency equation (2), which is the gap equation with $\bar{\lambda}$ as gap parameter Δ ($\equiv U m_s$), instead of three self-consistency equations (2)–(4).

The energy of the AF state (per site) and the spin stiffness ρ_s are given by:

$$e_{\text{AF}} = -2q_s \int_{\bar{\mu}}^{\infty} d\varepsilon \mathcal{N}(\varepsilon) \sqrt{\varepsilon^2 + \bar{\lambda}^2} + U d^2 + \bar{\lambda} q_s m_s \quad (8)$$

$$\rho_s = \frac{q_s}{4} \int_{\bar{\mu}}^{\infty} d\varepsilon \frac{\mathcal{N}(\varepsilon) \varepsilon^2}{(\varepsilon^2 + \bar{\lambda}^2)^{1/2}} - \frac{z_+^2 z_-^2}{q_s} \int_{\bar{\mu}}^{\infty} d\varepsilon \frac{\mathcal{N}_v(\varepsilon) \varepsilon^2}{(\varepsilon^2 + \bar{\lambda}^2)^{3/2}} \quad (9)$$

$\mathcal{N}_v(\varepsilon)$ is the weighted DOS, which, as well as the combination $z_+ z_-$, is specified in Ref.[12]. I will not repeat the expressions here since I will now restrict to half-filling ($\bar{\mu} = 0$, $n = 1$) for which case $z_+ z_-$ equals zero. For half-filling, the band-renormalization reduces to:

$$q_s(1, m_s, d) = \frac{2d^2 \left[\sqrt{1 - 2d^2 + m_s} + \sqrt{1 - 2d^2 - m_s} \right]^2}{1 - m_s^2}. \quad (10)$$

3 Large- U expansion in terms of moments

The above set of equations (2)–(5) can be solved numerically if the DOS for freely hopping electrons is given. For instance, in two dimensions $\mathcal{N}(\varepsilon)$ is known analytically and numerical solutions to (2)–(5) were obtained in Ref.[12]. Now I will derive solutions to (2)–(5) for $n = 1$ in the form of a systematic series expansion in $1/U$. This expansion is derived in two steps: (i) From the explicit form of the equations above (or from numerical solutions) it can be seen that large U corresponds to large $\bar{\lambda}$. Therefore, I first make a large- $\bar{\lambda}$ expansion for all quantities of interest. Subsequently, (ii) the $\bar{\lambda}$ -expansion for U (following from equation (4)) is inverted and substituted in the large- $\bar{\lambda}$ expansion for all the relevant quantities obtained before. In this way, one obtains the desired large- U expansion.

By expanding the square root in the denominator of the integrals in (2)–(5) for $\bar{\lambda} \gg \varepsilon$ one obtains a large- $\bar{\lambda}$ expansion in terms of *moments* M_n of the DOS:

$$M_n = \int_{-\infty}^{\infty} d\varepsilon \mathcal{N}(\varepsilon) \varepsilon^n. \quad (11)$$

After some work one arrives at the following expansions in $q \equiv 1/\bar{\lambda}$ [13]:

$$m_s = 1 - \frac{1}{2} M_2 q^2 + \frac{3}{8} M_4 q^4 - \frac{5}{16} M_6 q^6 + \frac{35}{128} M_8 q^8 + \mathcal{O}(q^{10}), \quad (12)$$

$$d = \frac{1}{2} \sqrt{M_2} q - \frac{4M_2^2 + 3M_4}{16\sqrt{M_2}} q^3 + \frac{\sqrt{M_2}}{16} \left[M_2^2 + \frac{13}{2} M_4 - \frac{9M_4^2}{16M_2^2} + \frac{5M_6}{2M_2} \right] q^5 \\ - \frac{1}{2048M_2^{5/2}} \left[192M_2^6 + 368M_2^4 M_4 + 364M_2^2 M_4^2 + 27M_4^3 + 736M_2^3 M_6 - 120M_2 M_4 M_6 + 280M_2^2 M_8 \right] q^7 + \mathcal{O}(q^9), \quad (13)$$

$$q_s = 1 - \frac{M_2}{4} q^2 - \left[\frac{M_2^2 - 7M_4}{16} \right] q^4 \\ - \left[\frac{9M_2^3 - 10M_2 M_4 + 12M_4^2/M_2 + 26M_6}{64} \right] q^6 + \mathcal{O}(q^8), \quad (14)$$

$$U = \frac{2}{q} + \frac{5}{2}M_2 q + \left[\frac{11M_2^2 - 21M_4}{8} \right] q^3 + \mathcal{O}(q^5) , \quad (15)$$

$$e_{\text{AF}} = -\frac{1}{2}M_2 q + \left[\frac{3M_2^2 + M_4}{8} \right] q^3 + \left[\frac{M_2^3 - 22M_2M_4 - 2M_6}{32} \right] q^5 + \mathcal{O}(q^7) , \quad (16)$$

$$\begin{aligned} \rho_s = & \frac{M_2}{8} q - \left[\frac{M_2^2 + 2M_4}{32} \right] q^3 + \left[\frac{-M_2^3 + 9M_2M_4 + 6M_6}{128} \right] q^5 \\ & - \left[\frac{9M_2^4 - 12M_2^2M_4 + 26M_4^2 + 32M_2M_6 + 20M_8}{512} \right] q^7 + \mathcal{O}(q^9) . \end{aligned} \quad (17)$$

For notational convenience the hopping integral t is taken equal to 1 in this section. For m_s , d , and q_s I have listed precisely the number of terms that is needed in order to obtain the number of terms (three) that is listed for U . For e_{AF} and ρ_s the terms are given that can be found using the listed terms for m_s , d , q_s , and U . Note that this implies that, to obtain the leading order in q for U , one already needs three terms for m_s and two terms for d .

Inverting the expansion for U gives:

$$q = 2p + 10M_2 p^3 + (122M_2^2 - 42M_4)p^5 + \mathcal{O}(p^7) , \quad (18)$$

where I have introduced the notation: $p \equiv 1/U$. Substituting this into the q -expansions, one obtains the following large- U expansions:

$$\begin{aligned} m_s = & 1 - 2M_2 p^2 - [20M_2^2 - 6M_4] p^4 \\ & - [294M_2^3 - 204M_2M_4 + 20M_6] p^6 + \mathcal{O}(p^8) , \end{aligned} \quad (19)$$

$$d^2 = M_2 p^2 + [6M_2^2 - 3M_4] p^4 + [75M_2^3 - 70M_2M_4 + 10M_6] p^6 + \mathcal{O}(p^8) , \quad (20)$$

$$\begin{aligned} q_s = & 1 - M_2 p^2 - [11M_2^2 - 7M_4] p^4 \\ & - [176M_2^3 - 192M_2M_4 + 12M_4^2/M_2 + 26M_6] p^6 + \mathcal{O}(p^8) , \end{aligned} \quad (21)$$

$$e_{\text{AF}} = -M_2 p - [2M_2^2 - M_4] p^3 - [15M_2^3 - 14M_2M_4 + 2M_6] p^5 + \mathcal{O}(p^7) , \quad (22)$$

$$\rho_s = \frac{M_2}{4} p + \frac{1}{2} [2M_2^2 - M_4] p^3 + \frac{3}{4} [15M_2^3 - 14M_2M_4 + 2M_6] p^5 + \mathcal{O}(p^7) . \quad (23)$$

A consistency check on the above results is that the following general identity still holds for the expansions above:

$$\frac{\partial e_{\text{AF}}}{\partial U} = d^2 . \quad (24)$$

Because in the SBMF at half-filling the spin stiffness is given by $-\frac{1}{8}$ times the average kinetic energy [12], we also have:

$$\rho_s = -\frac{1}{8} (e_{\text{AF}} - U d^2) . \quad (25)$$

Combining the latter formula with (24) it immediately follows that the expansion for ρ_s follows directly from the expansion for e_{AF} , as is also seen in the way the expansions are written above.

It is of some interest to compare the present result with results from the HFA and other expressions in the literature. It is far easier to derive the corresponding expression to (22) in the HFA since there is only one consistency equation (the gap equation) instead of three, and there is no band-renormalization function q_s which requires perturbative expanding. Following the same procedures as for SBMF, I find for the HFA:

$$e_{\text{AF}} = -M_2 p - [M_2^2 - M_4] p^3 - [4M_2^3 - 6M_2 M_4 + 2M_6] p^5 + \mathcal{O}(p^7). \quad (26)$$

Comparing with (22), one sees that HFA and SBMF give the same coefficient for the leading order contribution, but that there are differences for the higher order contributions. Note that also for the higher order contributions the term in the coefficient involving the highest moment agrees between HFA and SBMF. Since the moments are always positive, it is also clear that for large U the energy in the SBMF approximation is always lower than in the HFA.

In dimensions $D = 2$ and $D = 3$ (square and simple cubic lattices), the results can also be compared with a large- U expansion by Takahashi [14]. He derives a rigorous expansion (i.e. without making approximations) for the half-filled Hubbard model in terms of spin correlation functions of the $s = \frac{1}{2}$ Heisenberg model. These correlation functions are then evaluated using the Linear Spin Wave (LSW) approximation. Below the results of the latter approach are compared to those of HFA and SBMF for $D = 2$ and $D = 3$ (the necessary moments are easily evaluated and may for instance be found in the appendix):

	$D = 2$	$D = 3$	
HFA	$-4p + 20p^3 - 192p^5 + \dots$	$-6p + 54p^3 - 1344p^5 + \dots$	
SBMF	$-4p + 4p^3 + 256p^5 + \dots$	$-6p + 18p^3 + 600p^5 + \dots$	(27)
LSW	$-4.63p + 34.6p^3 + \dots$	$-6.58p + 65.6p^3 + \dots$	

It is clear that for large enough U the LSW expansion will give the lower energy. The leading order coefficient is proportional to the ground-state energy of the $s = \frac{1}{2}$ Heisenberg antiferromagnet, which is known to be very well approximated by LSW [15, 16]. Both the HFA and SBMF do not do better than the mean-field result for the leading order coefficient. In view of the fact that SBMF is an improvement over the HFA, it is somewhat remarkable that the HFA results for the next-to-leading order resemble the LSW results more than the SBMF results do. Also the different sign of the coefficient of p^5 between HFA and SBMF is notable. Further discussion of the range of validity of the expansion is given below.

To end this section a quantitative discussion of the large- U expansion is also given. In Table I, the ground-state energy of the AF phase at half-filling as obtained from the large- U expansion within SBMF ($e_{\text{AF}}^{\text{Uexp}}$) is compared to the “exact SBMF” result [$e_{\text{AF}}^{\text{SBMF}}$, found by solving (2)–(5) numerically] and to variational Monte Carlo results using an (antiferromagnetic) Gutzwiller wave function ($e_{\text{AF}}^{\text{GWVMC}}$, from Ref.[17]) for $D = 2$. For $D = 3$, only $e_{\text{AF}}^{\text{Uexp}}$ and $e_{\text{AF}}^{\text{GWVMC}}$ are compared. Note that the large- U expansion approximates the “exact SBMF” result very well for $U/t \geq 7$; it is expected that similar agreement is found for $D = 3$, where it is more involved

to compute $e_{\text{AF}}^{\text{SBMF}}$ since $\mathcal{N}(\varepsilon)$ is not a known analytical function as it is for $D = 2$ [18]. This is consistent with the fact that from the coefficients in the explicit expansions for $D = 2$ and $D = 3$, formula (27), one would estimate the series to converge well for $U/t \geq 8$ and $U/t \geq 6$ for two and three dimensions, respectively. SBFM is equivalent to the Gutzwiller *approximation* to the Gutzwiller wave function [9] and this approximation becomes exact in the limit of an infinite number of spatial dimensions [19, 20]. From the comparison with the GWVMC results it can be seen that the approximation is quite good already for $D = 2$. In view of the facts, that for $D = 3$ a very small lattice was used in the GWVMC calculations and a larger lattice will even raise the energy somewhat (the effect will be larger for smaller U), the agreement for $D = 3$ can even be called excellent. So the clear advantage of the large- U expansion is that it can give results that are, for $U/t \geq 8$, within 1% of the “exact SBFM” and GWVMC calculations (which especially for $D = 3$ are much more involved and computationally demanding) by means of the simple formula (22).

U	$D = 2$			$D = 3$	
	$-e_{\text{AF}}^{\text{SBMF}}$	$-e_{\text{AF}}^{\text{Uexp}}$	$-e_{\text{AF}}^{\text{GWVMC}}$	$-e_{\text{AF}}^{\text{Uexp}}$	$-e_{\text{AF}}^{\text{GWVMC}}$
6	0.623639	0.615226	0.629(3)	0.8395	0.886(5)
8	0.485104	0.484375	0.493(3)	0.6965	0.704(7)
10	0.393528	0.393440	0.401(4)	0.5760	0.579(6)
12	0.330002	0.329990	0.336(5)	0.4872	0.491(7)
16	0.248782	0.248779	-	0.3700	-
20	0.199420	0.199420	-	0.2976	-

Table I. Numerical results in $D = 2$ and $D = 3$ for the ground-state energy of the antiferromagnetic phase at half-filling. $e_{\text{AF}}^{\text{Uexp}}$ is evaluated using the large- U expansion (22) and $e_{\text{AF}}^{\text{SBMF}}$ is obtained from (8) after solving (2)–(5). The GWVMC results are taken from Ref.[17] and were obtained on 20×20 and $6 \times 6 \times 6$ lattices for $D = 2$ and $D = 3$, respectively. Between brackets the statistical error in the last digit is given.

4 Large- U expansion for arbitrary dimension

In the appendix, it is shown that the moments M_n can be obtained as a function of dimension D [formula (A.9)]. If one substitutes these expressions in (22) and (26) one obtains a simultaneous $1/U$ and $1/D$ expansion for the energy of the AF phase:

$$\text{SBFM} \quad e_{\text{AF}} = -\frac{2Dt^2}{U} + \left(1 - \frac{3}{2D}\right) \frac{4D^2t^4}{U^3} + \left(-3 + \frac{24}{D} - \frac{20}{D^2}\right) \frac{8D^3t^6}{U^5} + \dots (28)$$

$$\text{HFA} \quad e_{\text{AF}} = -\frac{2Dt^2}{U} + \left(2 - \frac{3}{2D}\right) \frac{4D^2t^4}{U^3} + \left(-16 + \frac{36}{D} - \frac{20}{D^2}\right) \frac{8D^3t^6}{U^5} + \dots (29)$$

where I have included the hopping parameter t again. The above formulas are well suited to discuss the limit of an infinite number of dimensions. This is of interest since, as noted above, SBMF is equivalent to the Gutzwiller approximation to the Gutzwiller wave function; an approximation which becomes exact in the limit of an infinite number of spatial dimensions. Thus the present expansion allows us to scrutinize the approximation explicitly for lower dimension. In order for the limit $D \rightarrow \infty$ to be meaningful, one has to introduce a scaled hopping parameter $t^* = t\sqrt{2D}$ [19]. In terms of this scaled parameter the above SBMF large- U expansion assumes the form:

$$\frac{e_{\text{AF}}}{t^*} = -\frac{t^*}{U} + \left(1 - \frac{3}{2D}\right) \left(\frac{t^*}{U}\right)^3 + \left(-3 + \frac{24}{D} - \frac{20}{D^2}\right) \left(\frac{t^*}{U}\right)^5 + \dots \quad (30)$$

In Table II, I evaluate (30) for $D = 2, 3, 4$, and ∞ .

U^*	$-e_{\text{AF}}^*(D = 2)$	$-e_{\text{AF}}^*(D = 3)$	$-e_{\text{AF}}^*(D = 4)$	$-e_{\text{AF}}^*(D = \infty)$
2	0.34375	0.35069	0.36719	0.46875
4	0.24219	0.23947	0.23853	0.23730
8	0.12439	0.12394	0.12373	0.12314
12	0.08317	0.08303	0.08296	0.08277
16	0.06244	0.06238	0.06235	0.06226
20	0.04997	0.04994	0.04992	0.04988

Table II. Scaled ground-state energy of the AF phase at half-filling $e_{\text{AF}}^* \equiv e_{\text{AF}}/t^*$ within SBMF as a function of scaled interaction $U^* \equiv U/t^*$ for various dimensions D (with $t^* = t\sqrt{2D}$).

From Table II, one sees that for scaled interaction $U^* \geq 4$ (which corresponds to $U/t = 8$ for $D = 2$) already the result for $D = 2$ and the infinite-dimensional result coincide remarkably well. The simultaneous large- U , large- D expansion shows in detail how well the Gutzwiller approximation, and therefore SBMF, reproduces the Gutzwiller wave function for the various dimensions. It would have been interesting to compare the coefficients in the present results with those obtained by Gebhard in his $1/D$ -expansion for the half-filled Hubbard model [20]. Unfortunately, a direct comparison is not meaningful, since Gebhard only considers the paramagnetic Gutzwiller wave function and does not treat the antiferromagnetic phase at half-filling in a $1/D$ -expansion.

5 Discussion and Conclusions

There is no fundamental reason why the large- U expansion given here cannot be extended to higher orders in t/U . With a little more effort (especially the coefficients in the large- $\bar{\lambda}$ expansion of d in formula (13) require some care) more terms can be obtained. However, as already remarked in Section 3, the present series reproduces well the “exact SBMF” result for $U/t \geq 7$ in two dimensions, and this agreement is expected to be even better in three dimensions. Also, for smaller values of U/t the contribution from the $(t/U)^5$ term becomes larger than that from the $(t/U)^3$ term, which is an indication that the intrinsic radius of convergence of the series is close to $U/t = 7$. Therefore, extending the series can not be considered very useful.

A more interesting extension of the present series is to go off half-filling, i.e. $n \neq 1$. Again there is no fundamental reason preventing this, but now the task is quite a bit more laborious for a number of reasons. Not only are we dealing with the case $\bar{\mu} \neq 0$ and consequently *partial* moments are required [see (11)], which can only be computed numerically, also the expression for the band renormalization q_s , formulae (6)–(7), needs to be expanded for $n \neq 1$, which implies that non-rational coefficients in the expansion in terms of (partial) moments appear. Furthermore, in formula (9) for ρ_s also the term with the combination $z_+ z_-$ needs to be taken into account and perturbatively expanded. If similar large- U expansions are possible for more complicated phases, like spirals or domain walls, this would allow for a study of aspects of the phase diagram off half-filling.

In conclusion, a large- U expansion is derived for the antiferromagnetic phase of the Hubbard model within the framework of the slave-boson mean-field approximation. Even though such an expansion constitutes in a sense an approximation of an approximation, the resulting analytic expression is capable of reproducing very well results of elaborate Monte Carlo calculations. Furthermore, the expansion allows for an explicit study of the limit of a large number of spatial dimensions, since the occurring moments of the non-interacting density of states are obtained for arbitrary dimension.

Acknowledgments

I acknowledge useful discussions with P.G.J. van Dongen, F. Gebhard, and J.M.J. van Leeuwen on the work reported here, as well as comments by D.P. Aalberts on an earlier version of the paper.

Appendix

In this appendix some general expressions are derived for the moments M_n of the density of states $\mathcal{N}(\varepsilon)$ of freely hopping electrons on hypercubic lattices:

$$M_n = \int_{-\infty}^{\infty} d\varepsilon \mathcal{N}(\varepsilon) \varepsilon^n = \frac{(2t)^n}{(2\pi)^D} \int_{-\pi}^{\pi} \cdots \int_{-\pi}^{\pi} dk_1 \dots dk_D [\cos k_1 + \cdots + \cos k_D]^n . \quad (\text{A.1})$$

Although such moments are frequently used and calculated in the literature I have never encountered the general analytic expressions and simple formulae to be given below. First, I will derive simple expressions for *all* moments in the *lower* dimensions $D = 1, 2$, and 3. Then I will derive expressions for the *lower* moments in *arbitrary* dimension D .

By employing the multinomial generalization of the binomial it is possible to further work out (A.1):

$$M_n = (2t)^n \sum'_{n_1, \dots, n_D=0}^n \frac{n!}{n_1! \dots n_D!} q_{n_1} \dots q_{n_D} , \quad (\text{A.2})$$

where the prime on the sum denotes the restriction: $n_1 + \cdots + n_D = n$ and

$$q_m = \frac{1}{2\pi} \int_0^{2\pi} dk \cos^m(k) = \begin{cases} \frac{(m-1)!!}{m!!} & m \text{ even} \\ 0 & m \text{ odd} \end{cases} \quad (\text{A.3})$$

From this general form it is clear that all moments with n odd are zero for any dimension. All formulae below will be for moments with n even. For $D = 1$ the restriction allows for only one term in the sum and the result is trivial:

$$M_n = (2t)^n \frac{(n-1)!!}{n!!} \quad D = 1 . \quad (\text{A.4})$$

A more remarkable result is that the moments for $D = 2$ are exactly the squares of the moments for $D = 1$ (if t is put equal to 1):

$$M_n = t^n \left[2^n \frac{(n-1)!!}{n!!} \right]^2 \quad D = 2 . \quad (\text{A.5})$$

The proof of (A.5) follows directly from (A.2) by some manipulations with the factorials and employing the identity [21]:

$$\sum_{n=0}^N \binom{N}{n}^2 = \binom{2N}{N} . \quad (\text{A.6})$$

For $D = 3$ I have not been able to obtain an explicit expression for M_n , but the triple sum in (A.2) can be reduced to one simple, unrestricted sum. Using the relations: $(m-1)!!/m! = 1/m!! = 2^{-m/2}/(m/2)!$ (m even), one can rewrite (A.2) (for n even) as:

$$M_n = n! \sum'_{u,v,w=0}^{n/2} \frac{1}{(u!)^2 (v!)^2 (w!)^2} , \quad (\text{A.7})$$

where the prime indicates the constraint: $u + v + w = n/2$. This expression was already given by Brinkman and Rice for the moments of the density of states of one hole in a ferromagnetic background [22]. It can however be simply reduced to a single sum by incorporating the constraint and a subsequent application of (A.6). The result is:

$$M_n = n! \sum_{u=0}^{n/2} \frac{(n-2u)!}{(u!)^2 ((n/2-u)!)^4} \quad D = 3. \quad (\text{A.8})$$

Formulas (A.5) and (A.8) make for much easier evaluation of the moments than is usually found in the literature: e.g. in Ref.[23] moments up to $n = 22$ and $n = 16$ are computed for $D = 2$ and $D = 3$, respectively, using the method of counting the number of paths that return to the origin after n steps. I note that also finding an explicit expression for $D = 3$ may be possible on account of the (empirical) fact that the moments for $D = 3$ always contain the moments for $D = 1$ as a factor.

The general expression (A.2) shows that finding the terms contributing to M_n amounts to finding all the combinations of D even numbers (including zero) that add up to n . For the lower moments the number of possibilities is not large and can be expressed in terms of D using combinatorial arguments. I have found the following results:

$$\begin{aligned} M_2 &= 2D \\ M_4 &= 6D(2D-1) \\ M_6 &= 20D(6D^2-9D+4) \\ M_8 &= 70D(24D^3-72D^2+82D-33) \\ M_{10} &= 252D(14400D^4-143400D^3+501050D^2-715225D+343176) \end{aligned} \quad (\text{A.9})$$

References

- [1] M.C. Gutzwiller, Phys. Rev. Lett. **10**, 159 (1963), J. Hubbard, Proc. Roy. Soc. (London) **A276**, 238 (1963), J. Kanamori, Progr. Theor. Phys. (Kyoto) **30**, 275 (1963), P.W. Anderson, Phys. Rev. **115**, 2 (1959).
- [2] M.S. Hybertsen, M. Schlüter, and N.E. Christensen, Phys. Rev. B **39**, 9028 (1989); G. A. Sawatzky, in *High-Temperature Superconductivity*, edited by D.P. Tunstall and W. Barford (Adam Hilger, Bristol, 1991).
- [3] J.R. Schrieffer, X.G. Wen, and S.C. Zhang, Phys. Rev. B **39**, 11663 (1989).
- [4] P. Nozières and S. Schmitt-Rink, J. Low Temp. Phys. **59**, 195 (1985), A.J. Leggett, in *Modern Trends in the Theory of Condensed Matter* (Springer, Berlin, 1980), p.13.
- [5] M. Randeria, J.-M. Duan, and L.-Y. Shieh, Phys. Rev. B **41**, 327 (1990).
- [6] P.J.H. Denteneer, Guozhong An, and J.M.J. van Leeuwen, Europhys. Lett. **16**, 5 (1991), Erratum, *ibid.* **16**, 509 (1991).

- [7] J. Carmelo and D. Baeriswyl, Phys. Rev. B **37**, 7541(1988).
- [8] G. Beni, T. Holstein, and P. Pincus, Phys. Rev. B **8**, 312 (1973).
- [9] G. Kotliar and A.E. Ruckenstein, Phys. Rev. Lett. **57**, 1362 (1986).
- [10] M.C. Gutzwiller, Phys. Rev. **134**, A923 (1964); Phys. Rev. **137**, A239 (1965).
- [11] The sign problem is more severe for large U/t (off half-filling) and most Quantum Monte Carlo work is for two dimensions (where $W = 8t$) and $U/t = 4$.
- [12] P.J.H. Denteneer and M. Blaauboer, J. Phys.: Condens. Matter **7**, 151 (1995), Erratum, *ibid.* **7**, 2377 (1995).
- [13] The expansion for m_s follows directly from (2), the expansion for d is first written with general coefficients which are then determined by the fact that (3) must be satisfied (this requires an expansion of $[q_s]_{m_s}$). The expansions for U , e_{AF} and ρ_s then follow from those for m_s and d (they require expansions of $[q_s]_d$ and q_s).
- [14] M. Takahashi, J. Phys. C **10**, 1289 (1977).
- [15] P.W. Anderson, Phys. Rev. **86**, 697 (1952).
- [16] D.C. Mattis, *The Theory of Magnetism* (Harper and Row, New York, 1982).
- [17] H. Yokoyama and H. Shiba, J. Phys. Soc. Jap. **56**, 3582 (1987).
- [18] The “exact SBMF” result for $D = 3$ is obtained by H. Hasegawa, Phys. Rev. B **41**, 9168 (1990); it is compared to the GWVMC result of Ref.[17] only graphically, but the quantitative agreement appears good.
- [19] W. Metzner and D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989).
- [20] F. Gebhard, Phys. Rev. B **41**, 9452 (1990).
- [21] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series, and Products*, formula 0.157 (Academic Press, New York, 1980).
- [22] W.F. Brinkman and T.M. Rice, Phys. Rev. B **2**, 1324 (1970).
- [23] D.F.B. ten Haaf, P.W. Brouwer, P.J.H. Denteneer, and J.M.J. van Leeuwen, Phys. Rev. B **51**, 353 (1995).